

Systems of Linear Equations4.1 Systems of Equations & Graphing (2x2)Objectives

- 1) Determine if an ordered pair is a solution of a system of linear equations.
- 2) Solve a system of linear equations using graphing
 - by hand
 - on GC (intersect)
- 3) Classify a system of linear equations
 - consistent independent
 - consistent dependent
 - inconsistentfrom the graph of the system.

4.2 Systems of Equations & Substitution (2x2)Objectives

- 1) Solve a system of linear equations using substitution.

4.3 Systems of Equations & Elimination (2x2)Objectives

- 1) Solve a system of linear equations using elimination.

GC 22: Intersection of Graphs

Quick ① Determine if $(-2.1, .3)$ is a solution of

$$\begin{cases} -\frac{x}{6} + \frac{y}{2} = \frac{1}{2} \\ \frac{x}{3} - \frac{y}{6} = -\frac{3}{4} \end{cases}$$

step 1: substitute $x = -2.1$ and $y = .3$ into 1st eqn (use GC!)

$$-(-2.1)/6 + .3/2$$

$$\boxed{\text{MATH}} \boxed{\text{ENTER}} \boxed{\text{ENTER}} \quad \text{>frac}$$

$$= \frac{1}{2} \checkmark$$

step 2: substitute $x = -2.1$ and $y = .3$ into 2nd eq'n (use GC!)

$$-2.1/3 - .3/6$$

$$\boxed{\text{MATH}} \boxed{\text{ENTER}} \boxed{\text{ENTER}}$$

$$= -3/4 \checkmark$$

step 3: Write yes or no

Yes $(-2.1, .3)$ makes both equations true.

Useful Observations:

1st: This system can be written with coefficients:

$$\begin{cases} -\frac{1}{6}x + \frac{1}{2}y = \frac{1}{2} \\ \frac{1}{3}x - \frac{1}{6}y = -\frac{3}{4} \end{cases}$$

2nd: We can clear fractions by multiplying each eq'n by its LCD.

$$\left(-\frac{x}{6} + \frac{y}{2} = \frac{1}{2}\right) \text{ by } 6 \Rightarrow -x + 3y = 3$$

$$\left(\frac{x}{3} - \frac{y}{6} = -\frac{3}{4}\right) \text{ by } 12 \Rightarrow 4x - 2y = -9$$

3rd: Soon we'll be able to solve this using matrices instead.

M70

$$\begin{cases} 1.5x + 2.5y = 22.05 & \text{(A)} \\ 3.2x - 0.7y = -15.9 & \text{(B)} \end{cases}$$

Solve by graphing on GC.

Approximate solution to nearest hundredth.

Step 1: Isolate y in each equation

$$\text{(A)} \quad 1.5x + 2.5y = 22.05$$

$$\frac{2.5y}{2.5} = \frac{-1.5x}{2.5} + \frac{22.05}{2.5}$$

$$y = -0.6x + 8.82$$

$$\text{(B)} \quad 3.2x - 0.7y = -15.9$$

$$\frac{-0.7y}{-0.7} = \frac{-3.2x}{-0.7} - \frac{15.9}{-0.7}$$

$$y = \frac{32}{7}x + \frac{159}{7}$$

Use **MATH**
>frac !!

* CAUTION * Do not use rounded decimals for equation (B) or your answers will include roundoff error


Step 2: Graph both lines in GC.

$$y_1 = -0.6x + 8.82$$

$$y_2 = 32/7 * x + 159/7$$

Step 3: Confirm visually that the point of intersection is visible in current GC window. If not, adjust the viewing window.

(YMAX must be MORE THAN 10)



Step 4: Use GC's **INTERSECT** command.

2nd **TRACE** = CALC

5 = Intersect

ENTER (1st curve) **ENTER** (2nd curve) **ENTER** (Guess)

② continued

Intersection

$$x = -2.68674 \quad y = 10.4322044$$

Step 5: Round each coordinate to the nearest hundredth and write an ordered pair.

$$\boxed{(-2.69, 10.43)}$$

③ Solve by substitution

* Review of Method

$$\begin{cases} 2.6x + y = 5.6 & \textcircled{A} \\ -4.3x - 2y = -4.9 & \textcircled{B} \end{cases}$$

Step 1: Identify best choice of variable to isolate
Isolate y in either equation before choosing x !

Solve $\textcircled{A}$ for y : $y = -2.6x + 5.6$

Step 2: Substitute into $\textcircled{B}$ to replace y and solve for x .

$$-4.3x - 2(-2.6x + 5.6) = -4.9$$

$$-4.3x + 5.2x - 11.2 = -4.9$$

$$.9x = 6.3$$

$$x = 7$$

Step 3: Substitute for x and solve for y .

$$y = -2.6(7) + 5.6$$

$$y = -12.6 = -\frac{63}{5}$$

Step 4: Write ordered pair

$$\boxed{(7, -12.6)} \text{ or } \boxed{(7, -\frac{63}{5})}$$

Note: Instructions did not say to round! Give exact answer.

what you will do!

④ Solve $\begin{cases} 3x - 2y = 10 & \textcircled{A} \\ 4x - 3y = 15 & \textcircled{B} \end{cases}$ by elimination.

Step 1: Identify best choice of variable to eliminate.

choice #1: eliminate x

$3x$ and $4x \Rightarrow$ the LCM of 3 and 4 is 12

mult $\textcircled{A}$ by 4: $3x \cdot 4 = 12x$

mult $\textcircled{B}$ by -3 : $4x \cdot (-3) = \frac{-12x}{0}$

★ choice #2: eliminate y

$-2y$ and $-3y \Rightarrow$ the LCM of 2 and 3 is 6

mult $\textcircled{A}$ by 3: $-2y \cdot 3 = -6y$

mult $\textcircled{B}$ by -2 : $-3y \cdot (-2) = \frac{6y}{0}$

Since 6 is smaller than 12, we choose option 2.

Step 2: Multiply all terms of each equation by multipliers needed to get LCM, with one equation positive and the other negative.

$\textcircled{A} \times 3$: $3 \cdot 3x - 3 \cdot 2y = 3 \cdot 10$
 $9x - 6y = 30$

$\textcircled{B} \times (-2)$: $-2 \cdot 4x - (-2) \cdot 3y = (-2) \cdot 15$
 $-8x + 6y = -30$

m70

④ cont

step 3: Add the two equations together, like terms to like terms. If necessary, isolate variable.

$$\begin{array}{rcl} 9x - 6y & = & 30 \quad \textcircled{A} \\ -8x + 6y & = & -30 \quad \textcircled{B} \\ \hline x & & = 0 \end{array}$$

step 4: Substitute result into any previous equation and solve for remaining variable.

subst $x=0$ into $\textcircled{A}$:

$$3(0) - 2y = 10$$

$$-2y = 10$$

$$y = -5$$

step 5: Write solution as an ordered pair.

$$\boxed{(0, -5)}$$

PREVIEW

Notice: It seems silly at this point, but we can write the solution as a system:

$$\begin{cases} x = 0 \\ y = -5 \end{cases}$$

or a more filled-in system

$$\begin{cases} 1x + 0y = 0 \\ 0x + 1y = -5 \end{cases}$$

Why? So we can use matrices on GC.

"Solve" vs "Classify"


Solve: means to find the values of the variables that make the system of equations true.

(I)

• The answer is an ordered pair

$$(a, b)$$

at the point where l_1 intersects l_2




OR

(II)

• There is no solution,

$$\emptyset$$

because l_1 and l_2 are parallel and do not intersect.



OR


(III)

• There are infinitely many solutions

$$\{(x, y) : \underline{\hspace{2cm}}\}$$

↑
write equation
of line here

because l_1 and l_2 are the same line when graphed.



Classify: means to identify the characteristics of the system in words.

{ consistent: the system has at least one solution.

{ inconsistent: the system has no solution

{ independent: none of the equations is a combination of the other equations


{ dependent: at least one equation is a combination of the other equations.

To classify a system (2×2), write two words:

(I) This system has a solution $\Rightarrow$
consistent

The two equations are different.


independent



(II) This system has no solution $\Rightarrow$
inconsistent

With 2×2 , we don't typically say that the equations are

independent




(III) Every point on one line is also on the other line—
 This system has an infinite number of solutions

consistent

But the two lines are the same line

dependent




When solving a consistent & dependent 2×2 system

CAUTION:

Infinitely many solutions (x, y)
is NOT "All real numbers x " as for a single equation.

System:



Solutions must be ON THE LINE to satisfy the equations.

• a real-number pair that is not on the line is not a solution. So "all real number pairs" do NOT work in the equations.

solution $\{(x, y) : \underline{\text{eqn}}\}$

Equation:



solutions on x -axis only.
Any x -value is all solution $\Rightarrow$ "all real numbers"

$\{x : x \in \mathbb{R}\}$

Math 70

⑦ Solve and classify.

$$\begin{cases} 3x + \frac{y}{2} = 2 & \textcircled{A} \\ 6x + y = 5 & \textcircled{B} \end{cases}$$

Method 1: Elimination

$$\begin{cases} 3x + \frac{y}{2} = 2 & \textcircled{A} \\ 6x + y = 5 & \textcircled{B} \end{cases} \leftarrow \begin{array}{l} \text{mult by } -2 \text{ to eliminate } x \\ \text{(or to eliminate } y) \end{array}$$

$$3(-2)x + \frac{y(-2)}{2} = 2(-2) \quad \textcircled{A} \times (-2)$$

$$\begin{cases} -6x - y = -4 & \textcircled{A}_{\text{new}} \\ 6x + y = 5 & \textcircled{B} \end{cases}$$

$$0x + 0x = 1$$

$$0 = 1$$

add equations
like terms w/ like terms

$$0 \neq 1 \text{ !?!}$$

False statement $\Rightarrow$ parallel lines

solve: no solution or $\emptyset$

classify: inconsistent or inconsistent independent

Method 2: substitution

$$\text{Solve } \textcircled{B} \text{ for } y: \quad \begin{array}{rcl} 6x + y & = & 5 \\ \underline{-6x} & & \underline{-6x} \end{array}$$

$$y = -6x + 5$$

$$\text{Substitute into } \textcircled{A}: \quad 3x + \frac{-6x + 5}{2} = 2$$

$$3x + \frac{-6x}{2} + \frac{5}{2} = 2$$

$\rightarrow$

$$\textcircled{7} \text{ cont } \underbrace{3x - 3x} + \frac{5}{2} = 2$$

$$0x + \frac{5}{2} = 2$$

$$\frac{5}{2} = 2$$

$\frac{5}{2} \neq 2$!?! false statement

Solve: no solution or ϕ

classify: inconsistent or inconsistent independent

* When using an algebraic or GC method and our correct work leads to a false statement containing only numbers (no variables), the system has no solution.

Math 70

(8) Solve and classify.

$$\begin{cases} y = \frac{1}{7}x + 3 & \textcircled{A} \\ x - 7y = -21 & \textcircled{B} \end{cases}$$

Method 1: Substitution

(A) already has y isolated.

Substitute (A) $\rightarrow$ (B)

$$x - 7\left(\frac{1}{7}x + 3\right) = -21$$

$$x - 7 \cdot \frac{1}{7}x - 7 \cdot 3 = -21$$

distribute

$$\underbrace{x - x} - 21 = -21$$

$$0x - 21 = -21$$

$$-21 = -21 \quad \text{true!?!}$$

solve:

$$\{(x, y) : x - 7y = -21\}$$

or

$$\{(x, y) : y = \frac{1}{7}x + 3\}$$

classify:

consistent
dependent

Caution: MathXL provides you with a completely filled-in set! But when you take a PQ or exam, you'll have to write it out yourself!!

Explore: What does $\{(x, y) : y = \frac{1}{7}x + 3\}$ mean?

"The set of all ordered pairs (x, y) so that $y = \frac{1}{7}x + 3$."

example: choose any value of x , plug into eqn.

$$x = 6 \Rightarrow y = \frac{1}{7}(6) + 3 = \frac{27}{7} \Rightarrow (6, \frac{27}{7}) \text{ is a solution.}$$

We can repeat for an infinite number of values of x to find many solutions.

⑧ cont

Method 2: Elimination

Equation (A) is not in standard form.

$$y = \frac{1}{7}x + 3 \quad (A)$$

subtract $\frac{1}{7}x$ both sides
to get $ax + by = c$

$$\begin{cases} -\frac{1}{7}x + y = 3 & (A) \\ x - 7y = -21 & (B) \end{cases}$$

$$-\frac{1}{7} \cdot 7x + 7 \cdot y = 3 \cdot 7 \quad (A) \times 7$$

$$\begin{cases} -x + 7y = 21 & (A) \\ x - 7y = -21 & (B) \end{cases}$$

To eliminate x , multiply equation (A) by 7.
OR To eliminate y , multiply equation (A) by 7.
OR To eliminate x , multiply equation (B) by $\frac{1}{7}$.
OR To eliminate y , multiply equation (B) by $\frac{1}{7}$.

$$0x + 0y = 0$$

add like
terms each side

$$0 = 0 \quad \text{true!?!}$$

solve: $\{(x, y) : x - 7y = -21\}$

or $\{(x, y) : y = \frac{1}{7}x + 3\}$

classify:

consistent
dependent

Extras:


- ① Solve by graphing on GC. Approximate to the nearest hundredth.

$$\begin{cases} y + 2.6x = 5.6 \\ y - 4.3x = -4.9 \end{cases}$$

$$\begin{cases} y_1 = -2.6x + 5.6 \\ y_2 = 4.3x - 4.9 \end{cases}$$

y=

ZOOM 6



2nd TRACE = CALC

5. Intersect

ENTER³

Intersection

$$X = 1.5217391$$

$$Y = 1.6434783$$

Round to nearest hundredth

(1.52, 1.64)

- 2 Solve the system of equations by the elimination method.

$$\begin{cases} \frac{5}{9}x + \frac{1}{3}y = \frac{34}{3} & \leftarrow \text{LCD} = 9 \\ \frac{4}{81}x - \frac{5}{9}y = -\frac{115}{27} & \leftarrow \text{LCD} = 81 \end{cases}$$

Select the correct choice below and, if necessary, fill in the answer box to complete your choice.

should be $\rightarrow$ ☐ A. The solution of the system is .

(Simplify your answer. Type an ordered pair. Use integers or fractions for any numbers in the expression.)

☐ B. There are infinitely many solutions.

☐ C. There is no solution.

clear fractions!

$$\begin{cases} \frac{5}{9}x \cdot 9 + \frac{1}{3}y \cdot 9 = \frac{34}{3} \cdot 9 \\ \frac{4}{81}x \cdot 81 - \frac{5}{9}y \cdot 81 = -\frac{115}{27} \cdot 81 \end{cases}$$

$$\begin{cases} 5x + 3y = 102 & \textcircled{A} \\ 4x - 45y = -345 & \textcircled{B} \end{cases}$$

elim y by mult $\textcircled{A}$ by 15

$$\begin{array}{r} 75x + 45y = 1530 \\ 4x - 45y = -345 \\ \hline \end{array}$$

$$79x = 1185$$

$$x = 15$$

Subst back to A

$$5(15) + 3y = 102$$

$$3y = 27$$

$$y = 9$$

$$\boxed{(15, 9)}$$

Way Easier: Use GC

MATRIX 2×3

$$\begin{bmatrix} 5/9 & 1/3 & 34/3 \\ 4/81 & -5/9 & -115/27 \end{bmatrix}$$

RREF

$$\begin{bmatrix} 1 & 0 & 15 \\ 0 & 1 & 9 \end{bmatrix}$$

$$\boxed{(15, 9)}$$