

## Objectives

1) Direct translation for numbers

"reciprocal of  $x$ " =  $\frac{1}{x}$

"quotient of  $x$  and 5" =  $\frac{x}{5}$

"sum" or "difference" means use parentheses or entire quantity above or below a fraction bar

2) Proportions

one ratio = another ratio

align like quantities down and across, not diagonally to set up the proportion

solve by cross-multiply

3) Work rates

- Find the fractions done by each worker in one unit of time

Total job = 5 hrs  $\Rightarrow \frac{1}{5}$  of job in one hour

- Add fractions for each worker

- Find fraction done by all workers in one unit of time

Total job =  $x$  min  $\Rightarrow \frac{1}{x}$  of job in one minute

- set sum of individual fractions = fraction together

4)  $D=RT$

- use a chart

- If you have  $D$  and  $R$ , use  $\frac{D}{R} = T$

- If "same time" given, set  $\frac{D_1}{R_1} = \frac{D_2}{R_2}$

- If you have  $D$  and  $T$ , use  $\frac{D}{T} = R$

- If "same rate" given, set  $\frac{D_1}{T_1} = \frac{D_2}{T_2}$

5) Random equation given in the problem.

- check what units go with each variable

- Use units on given numbers to identify which variable

## Objectives

- 1) Direct Translation
  - a. "quotient of a and b" means  $\frac{a}{b}$
  - b. "reciprocal of a" means  $\frac{1}{a}$
- 2) Proportions
  - a. Set up by checking vertically and horizontally
  - b. Solve by cross-multiplying
- 3) Work rates
- 4) Uniform Motion ( $D=RT$ )
- 5) Solve formulas given in the question

## Examples

Solve.

- 1) Five divided by the sum of a number and 4, minus the quotient of 3 and the difference of the number and 4 is equal to 6 times the reciprocal of the difference of the number squared and 16. What is the number?
- 2) To estimate the number of people in Springfield, population 10,000, who have a swimming pool in their backyard, 250 people were interviewed. Of those polled, 108 had a swimming pool. How many people in the city might one expect to have a swimming pool? (Round to the nearest whole number, if necessary.)
- 3) A painter can finish painting house in 5 hours. Her assistant takes 7 hours to finish the same job. How long would it take for them to complete the job if they were working together?
- 4) A baker can decorate the day's cookie supply four times as fast as his new assistant.
- 5) A car travels 400 miles on level terrain in the same amount of time it travels 160 miles on mountainous terrain. If the rate of the car is 30 miles per hour less in the mountains than on level ground, find its rate in the mountains.
- 6) In electronics, the relationship among the resistances  $R_1$  and  $R_2$  of two resistors wired in a parallel circuit and their combined resistance  $R$  is described by the formula  $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$ . If the combined resistance of two resistors wired in a parallel circuit is 2 ohms and one of the two resistances is 8 ohms, find the other resistance.

## Extra Problems

Solve.

- 7) One pump can drain a pool in 9 minutes. When a second pump is also used, the pool only takes 4 minutes to drain. How long would it take the second pump to drain the pool if it were the only pump in use?
- 8) A cyclist bikes at a constant speed for 16 miles. He then returns home at the same speed but takes a different route. His return trip takes one hour longer and is 21 miles. Find his speed.
- 9) One conveyor belt can move 1000 boxes in 7 minutes. Another can move 1000 boxes in 10 minutes. If another conveyor belt is added and all three are used, the boxes are moved in 3 minutes. How long would it take the third conveyor belt along to do the same job?
- 10) A boat moves 9 km upstream in the same amount of time it moves 19 km downstream. If the rate of the current is 8 km per hour, find the rate of the boat in still water.
- 11) A recent advertisement claimed that 2 out of every 3 doctors recommend a certain herbal supplement to increase energy levels. If a local hospital employs 250 doctors, how many doctors would you expect to recommend the supplement? (Round to the nearest whole number, if necessary.)
- 12) Jim can run 5 miles per hour on level ground on a still day. One windy day, he runs 15 miles with the wind, and in the same amount of time runs 4 miles against the wind. What is the rate of the wind?
- 13) One way to gauge whether a person may need to gain or lose weight is to use the body-mass index, which is calculated using the formula  $B = \frac{705w}{h^2}$ , where  $w$  is the weight in pounds and  $h$  is the height in inches. Doctors recommend that body-mass index values fall between 19 and 25. If a person is 6 ft 6 in. tall and has a body-mass index of 24, what should his or her weight be? (Round to the nearest pound.)
- 14) Five divided by the difference of a number and 8 equals the quotient of 10 and the sum of the number and 4. Find the number.
- 15) Two times the reciprocal of a number equals 28 times the reciprocal of 35. Find the number.
- 16) Mark and Rachel both work for Smith Landscaping Company. Mark can finish a planting job in 2 hours, while it takes Rachel 4 hours to finish the same job. If Mark and Rachel will work together on the job, and the cost of labor is \$40 per hour, what should the labor estimate be? (Round to the nearest cent, if necessary.)
- 17) In a race, Car A starts 1 mile behind Car B. Car A is traveling at 45 miles per hour, while Car B is traveling at 40 miles per hour. How long will it take for Car A to overtake Car B?

① Set up and solve - direct translate

$$\frac{5}{x+4} - \frac{3}{x-4} = 6 \cdot \frac{1}{x^2-16}$$

$$\frac{5}{x+4} - \frac{3}{x-4} = \frac{6}{(x+4)(x-4)}$$

domains:  $\begin{matrix} \uparrow & & \uparrow & & \uparrow \\ x \neq -4 & & x \neq 4 & & x \neq -4, 4 \end{matrix}$

$$\text{LED} = (x+4)(x-4)$$

$$\frac{5}{\cancel{(x+4)}} \cdot \cancel{(x+4)}(x-4) - \frac{3}{\cancel{(x-4)}} (x+4)\cancel{(x-4)} = \frac{6}{\cancel{(x+4)}\cancel{(x-4)}} \cdot \cancel{(x+4)}\cancel{(x-4)}$$

$$5(x-4) - 3(x+4) = 6$$

$$5x - 20 - 3x - 12 = 6$$

$$2x - 32 = 6$$

$$2x = 38$$

$$\boxed{x = 19}$$

② Proportion

$$\begin{array}{lcl} \text{pool/survey} \rightarrow \frac{108}{250} & = & \frac{x}{10,000} \leftarrow \text{pool/population} \\ \text{total/survey} \rightarrow & & \leftarrow \text{total/population} \end{array}$$

Check across and up & down that similar quantities are aligned.

Solve by cross-multiply:

$$(108)(10,000) = 250x$$

$$1080000 = 250x$$

$$\frac{1080000}{250} = x$$

$$4320$$

$$x = \boxed{4320 \text{ have pools}}$$

## (3) Setup and solve — work rate

painter: 5 hrs total job alone  $\Rightarrow \frac{1}{5}$  of job in one hour

assistant: 7 hrs total job alone  $\Rightarrow \frac{1}{7}$  of job in one hour

together:  $x$  hrs total  $\Rightarrow \frac{1}{x}$  of job in one hour

These fractions are called work rates.

Write equation using work rates, which all refer to one hour of work:

$$\frac{1}{5} + \frac{1}{7} = \frac{1}{x}$$

of job  
done  
by  
painter  
alone
of job  
done  
by  
assistant  
alone
of job  
done  
together

We expect  $x$  to be less than 5 because the painter alone takes 5 hrs and the assistant's help only shortens the time needed.

$$\text{LCD} = 35x$$

$$\frac{1}{5} \cdot \frac{35x}{35x} + \frac{1}{7} \cdot \frac{35x}{35x} = \frac{1}{x} \cdot \frac{35x}{35x}$$

$$7x + 5x = 35$$

$$12x = 35$$

$$x = \frac{35}{12} \text{ hrs}$$

$$x = \frac{35}{12} \text{ hrs} = \boxed{2\frac{11}{12} \text{ hrs}} = \boxed{2.91\bar{6} \text{ hrs}}$$

best answer

acceptable  
answer with  
all decimals  
and repeat bar

$$\boxed{\frac{35}{12} \text{ hrs}}$$

← ok, but  
harder to  
compare to 5  
and 7 to check  
if answer makes  
sense.

- ③ Method 2: Set up and solve — work rate  
Using units analysis to compare this question  
to a  $D=RT$  problem.

painter: 5 hrs per 1 job = rate  $\frac{5 \text{ hrs}}{1 \text{ job}}$  or  $\frac{1 \text{ job}}{5 \text{ hrs}}$

$$\begin{aligned} \text{amount of work done by painter} &= (\text{Rate}) \cdot (\text{Time}) \\ &= \frac{1 \text{ job}}{5 \text{ hrs}} \cdot t \text{ hrs} = \frac{t}{5} \text{ job} \end{aligned}$$

assistant: 7 hrs per 1 job = rate  $\frac{7 \text{ hrs}}{1 \text{ job}}$  or  $\frac{1 \text{ job}}{7 \text{ hrs}}$

$$\begin{aligned} \text{amount of work done by assistant} &= (\text{Rate}) \cdot (\text{Time}) \\ &= \frac{1 \text{ job}}{7 \text{ hrs}} \cdot t \text{ hrs} = \frac{t}{7} \text{ job} \end{aligned}$$

together:  $t$  hrs per 1 job = rate  $\frac{t \text{ hrs}}{1 \text{ job}}$  or  $\frac{1 \text{ job}}{t \text{ hrs}}$

$$\begin{aligned} \text{amount of work done together} &= (\text{Rate}) \cdot (\text{Time}) \\ &= \frac{1 \text{ job}}{t \text{ hrs}} \cdot t \text{ hrs} = \frac{t}{t} \text{ job} \\ &= 1 \text{ job} \end{aligned}$$

Equation:  $\left( \begin{array}{c} \text{work done} \\ \text{by painter} \end{array} \right) + \left( \begin{array}{c} \text{work done} \\ \text{by assistant} \end{array} \right) = \left( \begin{array}{c} \text{work done} \\ \text{together} \end{array} \right)$

$$\frac{t}{5} \text{ job} + \frac{t}{7} \text{ job} = 1 \text{ job}$$

$$\text{or } \frac{t}{5} + \frac{t}{7} = 1$$

$$\text{LCD} = 5 \cdot 7 = 35$$

$$35 \cdot \frac{t}{5} + 35 \cdot \frac{t}{7} = 35 \cdot 1$$

$$7t + 5t = 35$$

$$12t = 35$$

$$t = \boxed{\frac{35}{12} \text{ hrs}} \quad \text{same as Method 1.}$$

## ④ work rate

baker 4 times as fast as assistant.  
Whose work time (total) is shorter?

⇒ The baker's.

Let baker's work time =  $x$       fraction done in min  $\frac{1}{x}$   
assistant's work time =  $4x$       " " " "  $\frac{1}{4x}$

$$\frac{1}{x} + \frac{1}{4x} = \frac{1}{16}$$

baker      asst

solve for  $x$ , then find  $4x$  also.

↙ fractions done in one minute

$$\text{LCD} = 16x$$

$$16 + 4 = x$$

$$x = 20 \text{ min}$$

$$4x = 80 \text{ min}$$

less time ⇒ faster work

$x = \text{baker} = 20 \text{ mins}$ $4x = \text{asst} = 80 \text{ min}$
---

Caution: "faster" means "less time"

So faster person has a lower number.

$D = R \cdot T$  relates distance to rate (velocity or speed) and time.

$\frac{D}{R} = \frac{R \cdot T}{R}$  dividing both sides by  $R$  gives

$$\frac{D}{R} = T. \quad \text{Use this if "same time".}$$

$\frac{D}{T} = \frac{R \cdot T}{T}$  dividing both sides by  $T$  gives

$$\frac{D}{T} = R. \quad \text{Use this if "same rate".}$$

## ⑤

$$D = R \cdot T$$

$$\Rightarrow T = \frac{D}{R}$$

level	400	$R+30$	$\frac{400}{R}$
mtn	160	$R$	$\frac{160}{R-30}$

$$\frac{400}{R+30} = \frac{160}{R}$$

↑  
same time ⇒ use  $\frac{D}{R} = T$

↙ can also use  $R$   
 $R-30$

but  $R$  isn't the answer to the question!

⑤ cont LCD =  $R(R+30)$  or cross multiply

$$400R = 160(R+30)$$

$$400R = 160R + 4800$$

$$240R = 4800$$

$$R = 20 \text{ mph in mtns}$$

Method 2: 2 variables

	D	=	R	·	T
level	400		$R+30$		T
mtns	160		R		T

$$\begin{cases} 400 = T(R+30) \\ 160 = R \cdot T \end{cases}$$

$$T = \frac{160}{R}$$

$$400 = \frac{160}{R}(R+30)$$

$$400 = 160 + \frac{4800}{R}$$

$$240 = \frac{4800}{R}$$

$$240R = 4800$$

$$R = 20 \text{ mph}$$

Note: units analysis

$$(R+30) \frac{\text{mi}}{\text{hr}} \cdot (T) \text{ hr} = T(R+30) \text{ miles} = 400 \text{ miles}$$

$$(R) \frac{\text{mi}}{\text{hr}} \cdot (T) \text{ hr} = R \cdot T \text{ miles} = 160 \text{ miles}$$



⑥ Use given formula

$R$ ,  $R_1$ , and  $R_2$  are three different variables

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$$

$R$  = combined  
resistance  
given  
2

separate  
resistances

$R_1$  given (or  $R_2$  if you prefer)

← 3 variables in formula means  
2 numbers will be given so we  
can solve for one.

$$\frac{1}{2} = \frac{1}{8} + \frac{1}{R_1}$$

LCD of 2, 8, and  $R_1$  is  $8R_1$ .

$$\frac{1}{2} \cdot 8R_1 = \frac{1}{8} \cdot 8R_1 + \frac{1}{R_1} \cdot 8R_1$$

$$4R_1 = R_1 + 8$$

$$3R_1 = 8$$

$$R_1 = \frac{8}{3} \text{ ohms}$$

mult all by LCD

divide out/  
cancel common  
factors

subtract  $R_1$  both sides

⑦  $\frac{1}{9} + \frac{1}{x} = \frac{1}{4}$

1st pump fraction      2nd pump fraction      both pumps fraction

LCD of 9,  $x$ , and 4 is  $36x$

$$\frac{1}{9} \cdot 36x + \frac{1}{x} \cdot 36x = \frac{1}{4} \cdot 36x$$

mult all by LCD

$$4x + 36 = 9x$$

subtract  $4x$

$$36 = 5x$$

$$\frac{36}{5} = x$$

$$x = \frac{36}{5} \text{ min} = 7.2 \text{ min}$$

⑧  $D = R \cdot T$

16	$\frac{16}{T}$	T
21	$\frac{21}{T+1}$	T+1

↑

Same rate  $\Rightarrow R = \frac{D}{T}$

↓

Set rates equal

$$\frac{16}{T} = \frac{21}{T+1}$$

$$16(T+1) = 21(T)$$

$$16T + 16 = 21T$$

$$16 = 5T$$

time is  $3\frac{1}{5} \text{ hr} = \frac{16}{5} = T$

cross-multiply  
dist  
subtract  $16T$

But question asked for rate!

$$R = \frac{16}{T} = \frac{16}{(\frac{16}{5})} = 16 \div \frac{16}{5} = 16 \cdot \frac{5}{16} = \boxed{5 \frac{\text{mi}}{\text{hr}}}$$

Option 2:

$D = R \cdot T$

16	R	$\frac{16}{R}$	← T shorter time
21	R	$\frac{21}{R}$	← T+1 longer time

$$\frac{21}{R} = \frac{16}{R} + 1$$

$$\text{LCD} = R$$

to get longer time, add 1 to shorter time

$$21 = 16 + R$$

$$5 = R$$

$$\boxed{R = 5 \text{ mph}}$$

Option 3:

$D = R \cdot T$

16	R	T
21	R	T+1

$$\left. \begin{array}{l} 16 = R \cdot T \\ 21 = R(T+1) \end{array} \right\} \begin{array}{l} \text{solve system} \\ \text{by substitution} \end{array}$$

$R = \frac{16}{T}$  subst into  $21 = \frac{16}{T}(T+1)$  mult by T

$21T = 16(T+1)$  continue as before.

Math 70 6.6

⑨ "1000 boxes" = 1 job. The number 1000 is not needed.

$$\frac{1}{7} + \frac{1}{10} + \frac{1}{x} = \frac{1}{3}$$

fraction done by 1st      2nd      3rd      all 3 together

LCD of 7, 10, x, 3 = 210x

$$\frac{1}{7} \cdot 210x + \frac{1}{10} \cdot 210x + \frac{1}{x} \cdot 210x = \frac{1}{3} \cdot 210x$$

$$30x + 21x + 210 = 70x$$

$$51x + 210 = 70x$$

$$210 = 19x$$

$$\frac{210}{19} = x$$

$$x = \boxed{11\frac{1}{19} \text{ min}}$$

⑩ D = R · T

Going upstream is against the current

The current slows us down

RATE - CURRENT RATE  
(STILL WATER)

Going downstream is with the current

The current speeds us up.

RATE + CURRENT RATE  
(STILL WATER)

$$D = R \cdot T$$

upstream	9	$R - 8$	$\frac{9}{R-8}$
downstream	19	$R + 8$	$\frac{19}{R+8}$

↑  
same time

use  $\frac{D}{R} = T$

equation:  $\frac{9}{R-8} = \frac{19}{R+8}$

Math 70 6.6

(10) cont

LCD =  $(R-8)(R+8)$  or cross-multiply

$$9(R+8) = 19(R-8)$$

$$9R + 72 = 19R - 152$$

$$224 = 10R$$

$$\boxed{22.4 = R}$$

kph

(11)

$\frac{2}{3}$  of doctors

$\frac{2}{3}$  of 250

$$\frac{2}{3} \cdot 250 = \frac{500}{3} = 166.\overline{6}$$

round to the nearest whole doctor

$\boxed{167 \text{ doctors}}$

(12)

$$D = R \cdot T$$

Going against wind  
The wind slows us down

$$\text{RATE (no wind)} - \text{WIND RATE}$$

Going with wind  
The wind speeds us up

$$\text{RATE (no wind)} + \text{WIND RATE}$$

$$D = R \cdot T$$

against

with

4	$5-R$	$\frac{4}{5-R}$
15	$5+R$	$\frac{15}{5+R}$

↑  
same  
time

$$\text{use } \frac{D}{R} = T$$

equation:  $\frac{4}{5-R} = \frac{15}{5+R}$

LCD or cross-multiply

$$4(5+R) = 15(5-R)$$

$$20 + 4R = 75 - 15R$$

$$19R = 55$$

$$R = \frac{55}{19} = \boxed{2\frac{17}{19} \text{ mph}}$$

Math 70 6.6

(13) Use given formula

$$B = \frac{705W}{h^2}$$

$h$  = height in inches

height given 6 ft 6 in. =  $6 \cdot 12 + 6 = 78$  inches

$B$  = body mass index = 25

find weight.  $\Rightarrow$  solve for  $W$ .

$$24 = \frac{705W}{(78)^2}$$

$$24 = \frac{705W}{6084}$$

simplify  $78^2$

$$24(6084) = 705W$$

mult by 6084

$$146016 = 705W$$

$$\frac{146016}{705} = W$$

isolate  $W$ .

$$W = \frac{48672}{235}$$

> frac

$$W \approx 207.1148936$$

$$\boxed{W \approx 207 \text{ lbs}}$$

round to nearest pound

(14) Direct translation

$$\frac{5}{x-8} = \frac{10}{x+4}$$

LCD  $(x-8)(x+4)$

or cross-multiply

$$5(x+4) = 10(x-8)$$

$$5x + 20 = 10x - 80$$

dist

$$100 = 5x$$

$$\boxed{20 = x}$$

(15) Direct translation

$$2 \cdot \frac{1}{x} = 28 \cdot \frac{1}{35}$$

$$\frac{2}{x} = \frac{28}{35}$$

cross multiply

isolate

reduce

$$70 = 28x$$

$$x = \frac{70}{28} = \boxed{\frac{5}{2}}$$

(16)

$$\frac{1}{2} + \frac{1}{4} = \frac{1}{x}$$

Mark Rachel Together

$$\text{LCD} = 4x$$

$$\frac{1}{2} \cdot 4x + \frac{1}{4} \cdot 4x = \frac{1}{x} \cdot 4x$$

$$2x + x = 4$$

$$3x = 4$$

$$x = \frac{4}{3} \text{ hr} = 1\frac{1}{3} \text{ hr}$$

← fractions done  
in one hour.


Labor = \$40 per hour per person  $\Rightarrow$  \$80 per hour for both people

$$\text{Cost of labor for } \frac{4}{3} \text{ hr} = \frac{4}{3} \times 80 = \frac{320}{3} \approx \boxed{\$106.67}$$

(17)

$$D = R \cdot T$$

A	45T	45	T
B	40T	40	T



Same direction  $\Rightarrow$  subtract distances  
(not a rational equation)

$$45T - 40T = 1 \text{ mi}$$

$$5T = 1$$

$$\boxed{T = \frac{1}{5} \text{ hr}}$$

$$\frac{1}{5} \text{ hr} \cdot \frac{60 \text{ min}}{1 \text{ hr}} = \boxed{12 \text{ min}}$$

## Math 70 6.6

Extra A cyclist bikes at a constant speed for 23 miles. He returns home at the same speed but takes a different route. His return trip takes one hour longer and is 28 miles. Find his speed.

Alternate approach to the one in your class notes:  
(see 6.8 handout)

$$D = R \cdot T$$

23	$\frac{23}{T}$	T
28	$\frac{28}{T+1}$	T+1

Many of you wanted to use variables for time instead of for rate.

"Same speed"  
means:

$$\frac{23}{T} = \frac{28}{T+1}$$

Cross multiply

$$23(T+1) = 28T$$

$$23T + 23 = 28T$$

$$23 = 5T$$

$$4.6 = 4\frac{3}{5} = \frac{23}{5} = T$$

hrs      hrs      hrs

← This is the time to travel the 23 miles, not answer to question.

$$R = \frac{23}{T}$$

$$= \frac{23}{4.6}$$

$$R = 5 \text{ mph}$$

← To get speed, must plug back in.