

Objectives



- 1) Solve applications using variation
  - direct
  - inverse
  - joint

## Math 70 7.8 Variation

Objectives:

- 1) Direct variation
- 2) Inverse variation
- 3) Joint variation

Key words for recognizing variation problems: “varies” or “proportional”

|                | Model Words  | Translation to math   | Model equation  | Graph of model if $k = 2$  |
|----------------|--|---|---|--|
| <b>Direct</b>  | “y varies directly as x”<br><br>“y is directly proportional to x”<br><br>“y varies as x”   | “y varies” is always $y = k \cdot$<br><br>“directly” means x is in the numerator of RHS   | $y = kx$  |   |
| <b>Inverse</b> | “y varies inversely as x”<br><br>“y is inversely proportional to x”  | “y varies” is always $y = k \cdot$<br><br>“inversely” means x is in the denominator of RHS  | $y = k \cdot \frac{1}{x}$<br>or<br>$y = \frac{k}{x}$  |  |
| <b>Joint</b>   | “Joint” means that 3 or more variables are involved.<br><br><u>Example:</u> “y varies jointly as x and the square of w, and inversely as z and the cube root of v” | “y varies” is always $y = k \cdot$<br><br>“directly” means x and $w^2$ are in the numerator of RHS<br><br>“inversely” means z and $\sqrt[3]{w}$ are in the denominator of RHS | $y = k \cdot \frac{x \cdot w^2}{z \cdot \sqrt[3]{v}}$ | Not a 2-dimensional graph  |

Cautions:

- Direct variation problems can be solved by proportions. Inverse and Joint variation cannot!
- Variation equations have only multiply and divide, never add or subtract.
- Always write units on the final answer.
- Use units to help identify which values go with which variables.

**Process that works for ANY type of variation problem:**

Step 1: Define any variables, if needed. (Use letters that make sense.)

Step 2: Translate the sentence into an equation of variation. Don't forget k!

Step 3: Substitute a complete set of numbers (given in question) and solve for k.

Step 4: Solve the incomplete set of numbers (given in question) and solve to answer the question.

## Examples

1. Hooke's law states that the distance a spring stretches is directly proportional to the weight attached to the spring. If a 40-pound weight attached to a spring stretches the spring 5 inches, find the distance that a 65 pound weight will stretch that same spring.
2. Boyle's law says that if the temperature stays the same, the pressure  $P$  of a gas is inversely proportional to the volume  $V$ . If a cylinder in a steam engine has a pressure of 960 kilopascals when the volume is 1.4 cubic meters, find the pressure when the volume increases to 2.5 cubic meters.
3. The lateral surface area of a cylinder varies jointly as its radius and height.
  - a. Express this surface area  $S$  in terms of radius  $r$  and height  $h$ .
  - b. If the lateral surface area is  $20\pi$  square cm when the radius is 2 cm and the height is 5 cm, find the exact constant of variation and the equation of variation.
  - c. Find the radius when the lateral surface area is  $40\pi$  square cm and the height is 2 cm.
4. The maximum weight that a circular column can support is directly proportional to the fourth power of its diameter and is inversely proportional to the square of its height. A 2-meter-diameter column that is 8 meters in height can support 1 ton. Find the weight that a 1-meter-diameter column that is 4 meters in height can support.

Extras:

1. The maximum weight that a rectangular beam can support varies jointly as its width and the square of its height and inversely as its length. If a beam  $\frac{1}{2}$  foot wide,  $\frac{1}{3}$  foot high, and 10 feet long can support 12 tons, find how much a similar beam can support if the beam is  $\frac{2}{3}$  foot wide,  $\frac{1}{2}$  foot high, and 16 feet long.
2. The horsepower to drive a boat varies directly as the cube of the speed of the boat. If the speed of the boat is to double, determine the corresponding increase in horsepower required.
3. The volume of a cone varies jointly as its height and the square of its radius. If the volume of a cone is  $32\pi$  cubic inches when the radius is 4 inches and the height is 6 inches, find the volume of a cone when the radius is 3 inches and the height is 5 inches.
4. The intensity of light (in foot-candles) varies inversely as the square of  $x$ , the distance in feet from the light source. The intensity of light 2 feet from the source is 80 foot-candles. How far away is the source if the intensity of light is 5 foot-candles?

## Variation and Problem Solving

1) Direct variation.

2) Inverse variation

3) Joint variation

} Do not solve by proportions!

Key words for recognizing these problems are

\* varies

\* proportional.

However, "proportional" suggests that these problems should be solved using proportions, but only the first type can be solved using ordinary proportions.

Goal: one method for all 3 types of variation

First step: Translate each type of variation to an equation of variation

Direct Variation

model problem:  $y$  varies directly as  $x$

or:  $y$  is directly proportional to  $x$

mean  $y = k \cdot x$   $\leftarrow x$  in numerator of RHS

As  $x$  increases,  $y$  increases

$k$  is a number, a constant, called the constant of variation.

All variation problems have  $k$ .

$k$  is essential. The problem can't be done without it.

Inverse Variation

model problem:  $y$  varies inversely as  $x$

or:  $y$  is inversely proportional to  $x$ .

mean  $y = k \cdot \frac{1}{x}$   $\leftarrow x$  in denominator of RHS

or  $y = \frac{k}{x}$

As  $x$  increases,  $y$  decreases

Joint variation means

- 3 or more variables involved (not including  $k$ , which is not a variable)
- Each variable is either direct (in numerator) or inverse (in denominator).
- If problem does not say "direct" or "inverse", assume it's direct and write it in the numerator.

Examples of joint variation equations.

①  $y$  varies jointly as  $x$  and  $z$

means  $y = k \cdot x \cdot z$

②  $y$  varies jointly as  $x$  and inversely as  $z$

means  $y = k \cdot \frac{x}{z}$

③  $P$  varies jointly as  $q$  and the square of  $r$

means  $P = \frac{k \cdot q}{r^2}$

### Process for Solving Variation Problems

step 1: Translate to an equation of variation, with  $k$ .

step 2: Solve for  $k$ .

In the problem, you will be given a complete set of numbers, one for each variable.  
Plug them all in, and solve the result for  $k$ .

\* Once we know the value of  $k$ , we can use it for all of the rest of the problem. \*

step 3: Answer the question.

In the problem, you will be given a partial set of numbers, one for every variable except the question.  
Plug them all in, plug in the value of  $k$  from step 2, and solve for the requested variable.

Helpful notes:

- Variation equations have only multiply and divide.  
There is never add or subtract.

- Always write units on answers.

- Use units to help identify which values are which variables.

- Use letters that remind you of their meanings, especially in joint problems with many variables.

# Math 70 Practice Problems for Variation

1. Hooke's law states that the distance a spring stretches is directly proportional to the weight attached to the spring. If a 40-pound weight attached to a spring stretches the spring 5 inches, find the distance that a 65 pound weight will stretch that same spring.

step 1:  $D = k \cdot W$   $D = \text{distance}$   
 $W = \text{weight}$

step 2:  $5 = k \cdot 40$   
 $\frac{1}{8} = k$

step 3:  $D = \frac{1}{8} \cdot 65 = \boxed{\frac{65}{8} \text{ cm}} = \boxed{8.125 \text{ cm}}$

2. Boyle's law says that if the temperature stays the same, the pressure  $P$  of a gas is inversely proportional to the volume  $V$ . If a cylinder in a steam engine has a pressure of 960 kilopascals when the volume is 1.4 cubic meters, find the pressure when the volume increases to 2.5 cubic meters.

step 1:  $P = k \cdot \frac{1}{V}$

step 2:  $960 = \frac{k}{1.4}$

$1344 = k$

step 3:  $P = \frac{1344}{2.5}$

$P = \boxed{537.6 \text{ kilopascals}}$

3. The lateral surface area of a cylinder varies jointly as its radius and height.

a. Express this surface area  $S$  in terms of radius  $r$  and height  $h$ .

b. If the lateral surface area is  $20\pi$  square cm when the radius is 2 cm and the height is 5 cm, find the exact constant of variation and the equation of variation.

c. Find the radius when the lateral surface area is  $40\pi$  square cm and the height is 2 cm.

a = step 1:  $S = k \cdot r \cdot h$

b = step 2:  $20\pi = k \cdot 2 \cdot 5$   
 $2\pi = k$

rewrite eqn  $S = 2\pi r h$

c = step 3:  $40\pi = 2\pi \cdot r \cdot 2$   
 $10 \text{ cm} = r$

4. The maximum weight that a circular column can support is directly proportional to the fourth power of its diameter and is inversely proportional to the square of its height. A 2-meter-diameter column that is 8 meters in height can support 1 ton. Find the weight that a 1-meter-diameter column that is 4 meters in height can support.

max weight  
column  
supports

is directly  
proportional

4th power diameter  
inversely square of height

weight =  $W$   
diameter =  $d$   
height =  $h$

step 1:  $W = k \cdot \frac{d^4}{h^2}$

step 2:  $\left. \begin{matrix} d=2 \\ h=8 \\ W=1 \end{matrix} \right\} \Rightarrow 1 = \frac{k \cdot 2^4}{8^2} \Rightarrow 1 = \frac{16k}{64} \Rightarrow 64 = 16k \Rightarrow k = 4$

step 3:  $\left. \begin{matrix} W=? \\ d=1 \end{matrix} \right\} \Rightarrow W = \frac{4 \cdot 1^4}{4^2} \Rightarrow \boxed{W = \frac{1}{4} \text{ ton}}$

## Extras:

1. The maximum weight that a rectangular beam can support varies jointly as its width and the square of its height and inversely as its length. If a beam  $\frac{1}{2}$  foot wide,  $\frac{1}{3}$  foot high, and 10 feet long can support 12 tons, find how much a similar beam can support if the beam is  $\frac{2}{3}$  foot wide,  $\frac{1}{2}$  foot high, and 16 feet long.

step 1 max weight rectangular beam vary jointly -width -square height -inversely length

$$M = k \cdot \frac{w \cdot h^2}{l}$$

$M$  = max weight  
 $w$  = width  
 $h$  = height  
 $l$  = length

step 2:  $w = \frac{1}{2}$   
 $h = \frac{1}{3}$   
 $l = 10$   
 $M = 12$

$$12 = \frac{k \cdot (\frac{1}{2}) (\frac{1}{3})^2}{10} \Rightarrow 120 = \frac{1}{18} k \Rightarrow k = 2160$$

step 3:  $w = \frac{2}{3}$   
 $h = \frac{1}{2}$   
 $l = 16$   
 $M = ?$

$$M = \frac{2160 \cdot (\frac{2}{3}) \cdot (\frac{1}{2})^2}{16}$$

$M = 22.5 \text{ tons}$

2. The horsepower to drive a boat varies directly as the cube of the speed of the boat. If the speed of the boat is to double, determine the corresponding increase in horsepower required.

step 1: horsepower varies directly cube of speed  
 $H$  = horsepower  
 $s$  = speed

$$H = k \cdot s^3$$

step 2: double the speed  $\Rightarrow$  (2s) substitute and simplify

$$H = k (2s)^3$$

$$H = k \cdot 8 \cdot s^3$$

$$H = 8k \cdot s^3$$

horsepower must be multiplied by 8

3. The volume of a cone varies jointly as its height and the square of its radius. If the volume of a cone is  $32\pi$  cubic inches when the radius is 4 inches and the height is 6 inches, find the volume of a cone when the radius is 3 inches and the height is 5 inches.

step 1: volume varies jointly height square radius

$V$  = volume  
 $h$  = height  
 $r$  = radius

$$V = k \cdot h r^2$$

step 2:  $V = 32\pi$   
 $r = 4$   
 $h = 6$

$$32\pi = k \cdot 6 \cdot 4^2 \Rightarrow 32\pi = 96k \Rightarrow k = \frac{32\pi}{96} = \frac{\pi}{3}$$

step 3:  $V = ?$   
 $r = 3$   
 $h = 5$

$$V = \frac{\pi}{3} \cdot 5 \cdot 3^2 \Rightarrow V = 15\pi \text{ cubic inches} \text{ or } 15\pi \text{ in}^3$$

4. The intensity of light (in foot-candles) varies inversely as the square of  $x$ , the distance in feet from the light source. The intensity of light 2 feet from the source is 80 foot-candles. How far away is the source if the intensity of light is 5 foot-candles?

step 1: Intensity varies inversely square of  $x$  = distance

$I$  = intensity in foot-candles  
 $x$  = distance in feet

$$I = \frac{k}{x^2}$$

step 2:  $I = 80 \text{ ft-candles}$   
 $x = 2 \text{ ft}$

$$80 = \frac{k}{2^2} \Rightarrow 80 = \frac{k}{4} \Rightarrow k = 320$$

step 3:  $x = ?$   
 $I = 5$

$$5 = \frac{320}{x^2} \Rightarrow 5x^2 = 320 \Rightarrow x^2 = \frac{320}{5} \Rightarrow x^2 = 64$$

distance  $x$  cannot be negative  $\Rightarrow x = \pm \sqrt{64} \Rightarrow x = 8 \text{ ft}$